

Fall Prevention Conference, Honolulu, Hawaii 2007



The Role of Physical Activity in the Prevention of Falls in Older Age.

Debra J. Rose, PhD
Co-Director
Center for Successful Aging
California State University, Fullerton
and
Fall Prevention Center of Excellence
University of Southern California



Introduction



- Falls are costly for the individual and society
- Regular participation in physical activity is integral to good health and independence. Also lowers risk for falls and fall-related injuries.
- Most promising fall prevention strategies include physical activity or exercise



Falls Are Preventable



- ***Many identifiable risk factors***

- ***Muscle Weakness****
- ***Balance/gait problems****
- ***Prior fall***
- ***Vision****
- ***ADL limitations****
- ***Depression*/dementia***
- ***Medications***

- ***Intervention programs work!***

- ***Evidence shows 20-50% ↓ fall rates***
- ***“3 E’s”: Evaluation, **Exercise**, Environment***

- ***Systematic approach needed***



Roles for Physical Activity

- **Primary**


Prevent onset of pathology and system impairments

- **Secondary**

Slow progression of disease and system impairments

- **Tertiary**

Restoration of function to level that allows for independence in performance of daily activities



Benefits of Physical Activity in Reducing Falls

- Effective in improving balance and/or reducing fall incidence rates
 - Different settings (community, home)
 - Varied durations, intensity, frequency, and type of exercise
 - Different levels of risk
 - Group versus individually tailored home exercise programs
 - Type of provider



Effective Interventions include:

- Multicomponent exercise programs
- Gait, balance, motor coordination, and functional tasks
- Tai Chi
- Dance
- Walking

BUT not across all levels of risk



Why are the Research Findings Mixed?

■ Methodological Weaknesses

- Poor design and/or treatment of data
- Small sample sizes
- Insufficient information (e.g., randomization, blinding, intervention content)
- Limited follow-up data
- Non-standardized outcome measures and timing of follow-up assessments
- Varied dosage (frequency, intensity, duration)
- Little or no information about adherence, adverse events.
- Outcome measures often incongruous with type of intervention and/or limit generalization to daily function.


Translating Research into Practice

- Role and type of physical activity may differ according to level of fall risk
- Specificity, intensity, frequency, and duration of intervention needs to reflect level of fall risk
 - Well-rounded programs
 - Individually tailored programs
- Physical activity programs may need to include behavioral component for long-term participation



Translating Research into Practice

- In 2005, only 48.1% of older Americans met recommended PA guidelines
- Physical inactivity levels even higher among older adults with disability (56% report no leisure time activity).
- Factors that influence initiation and long-term participation vary by race, gender, level of disability, age.




Current Recommendations?

- Any type of moderate intensity activity that results in small increases in heart rate.
 - At least 30 minutes per day
 - At least 5 days per week
- OR
- Vigorous Activity that results in large increases in HR
 - At least 20 minutes per day
 - At least 3 days per week



Core Ingredients?

- Whole body functional activities that focus on improving:
 - ✓ Muscular endurance
 - ✓ Muscular strength and power
 - ✓ Aerobic endurance
 - ✓ Flexibility



Core Ingredients?

- Activities that stimulate multiple dimensions of balance:
 - ✓ Processing and integration of sensory information
 - ✓ Anticipatory and reactive control of action
 - ✓ Allocation of attention
 - ✓ Multidirectional and segmental coordination activities



Low-Risk: Physical Activity Serves Primary Role in Preventing Onset of Disability

- Many activity choices available
- Simplest and least resource intensive is walking.
- Acceptable among ethnically diverse groups.
- Walking poles can be used to increase energy expenditure or provide increased stability.



Moderate Risk: Physical Activity Serves Secondary Role in Slowing Progression of Disease/Impairment

- Benefit from programs more tailored to individual needs
- Specific balance and gait activities selected on basis of comprehensive evaluation of system impairments



High Risk: Physical Activity Serves Tertiary Role by Raising Physical Capacity to Perform BADLs

- Carefully tailored exercise program as part of a multifactorial intervention strategy
- Emphasis on building strength and endurance levels in a seated or supported standing position until balance and gait activities can be included.



Future Research Directions

- Lingering questions remain to be answered:
 - ✓ What type(s) of physical activity is most effective in reducing falls?
 - ✓ How does level of fall risk influence type and intensity of intervention strategy?
 - ✓ What type(s) of intervention strategies are effective in promoting long-term participation in physical activity and fall prevention activities across fall risk levels?



Future Research Directions

- ✓ Should a reduction in falls constitute the only outcome used to judge effectiveness?
- ✓ How do factors of ethnicity, socioeconomic status, and geographical location affect the type of physical activity intervention implemented?
- ✓ Is exercise effective in lowering fall risk among physically frail or cognitively impaired?



Summary

- Physical activity has an important role to play in preventing or lowering risk for falls in most settings.
- The specific type, intensity, duration, and frequency of physical activity selected is influenced by level of fall risk.
- Including a behavior-change component in any physical activity program will be critical for long-term participation in fall prevention activities.

Mahalo!



Selected References

- Barnett et al. Community-based exercise improves balance and reduces falls in at-risk older people. (2002); Austr J Phys Ther: 48(3):215-9.
- Belza B, Walwick J, Shiu-Thornton S, Schwartz S, Taylor M, LoGerfo J. Older adult perspectives on physical activity and exercise: Voices from multiple cultures. Preventing Chronic Disease (serial outline) 2004 Oct (date cited). Available from: URL: http://www.cdc.gov/ped/issues/2004/oct/04_0028.htm.
- Buchner DM, Cress ME, de Lateur BJ, Esselman PC, Margherita AJ, Price R, Wagner EH. The effect of strength and endurance training on gait, balance, fall risk, and health services use in community-living older adults. J Gerontol A Biol Sci Med Sci 1997; 52A(4): M218-24.
- Campbell AJ, Robertson MC. Implementation of multifactorial interventions for fall and fracture prevention. Age Ageing 2006; 35-S2: ii60-ii64.
- Close J, Ellis M, Hooper R, Glucksman E, Jackson S, Swift C. Prevention of falls in the elderly trial (PROFET): A randomised controlled trial. Lancet. 1999; 353: 93-7.
- Freedman VA, Hodgson N, Lynn J, Spillman BC, Waidmann T, Wilkinson AM et al. Promoting declines in the prevalence of late-life disability: Comparisons of three potentially high-impact interventions. Millbank Quarterly 2006; 84 (3): 493-520.
- Gillespie LD, Gillespie WJ, Robertson MC, Lamb SE, Cumming RG, Rowe BH. Interventions for preventing falls in elderly people. Cochrane Database Syst Rev 2003(4):CD000340.
- Lord SR, Ward JA, Williams P, Strudwick M. The effect of a 12-month exercise trial on balance, strength, and falls in older women: A randomized controlled trial. J Am Geriatr Soc 1995; 43:1198-1206.

Selected References

- Lord SR, Castell S, Corcoran J, Dayhew J, Matters B, Shan A, Williams P. The effect of group exercise on physical functioning and falls in frail older people living in retirement villages: A randomized, controlled trial. *J Am Geriatr Soc* 2003; 51: 1685-1692
- McAuley E, Jerome GJ, Elavsky S, Marquez DX, Ramsey SN. Predicting long-term maintenance of physical activity in older adults. *Prev Med* 2003; 37: 110-118.
- Rose DJ. Promoting functional independence in older adults at risk for falls: The need for a multidimensional programming approach. *J Aging Phys Activity* 2002; 10: 1-19.
- Rose DJ. Fallproof. A comprehensive balance and mobility program. Champaign (IL): Human Kinetics; 2003.
- Rose DJ. The Role of Physical Activity in the Prevention of Falls in Older Age 2007. Commissioned paper for World Health Organization. Can be accessed at:
http://www.who.int/ageing/projects/falls_prevention_older_age/en/index.html
- Schneider JK, Eveker A, Bronder DR, Meiner SE, Binder EF. Exercise training for older adults: Incentives and disincentives for participation. *J Gerontol Nursing* 2003; 29: 21-31.